

1/31

Fig. 1 A.





Fig. 1 B.



2/31

Fig.2.







Fig.3A.



4/31



5/31

Fig.4A.





Fig.4B.



6/31

Fig.4C.






Fig.4D.





Notes:

- This Strategy can be applied to reformatting of an individual Fab and to a mixture of different clones.
- Asc1*: For technical reasons the internal fragment contains as "Asc1-compatible" Mlu1 site; Asc1 and Mlu1 recognition sequences are destroyed.

8/31

Fig.6A.






Fig.6B.



9/31

Fig.6C.




10/31

Fig.7A.


PH1IgG

on bB SA-streptavidin-human bNTD/CTD

**Fig.7B.**

M 28B2 IgG

on bB SA-streptavidin-human bNTD/CTD



11/31

Fig.8A.






Fig.8B.




12/31

Fig.9A.

M27E11 IgG (with competition)
on bCTD, detection a-myc and r-a-mHRP

**Fig.9B.**


M28B2 IgG (with competition)
on bCTD, detection a-myc and r-a-mHRP




13/31

Fig.9C.

Control antibodies (with competition)
on bCTD, detection anti-mouse or anti-goat HRP



**Fig.9D.**

PH1 IgG (with competition)
on bCTD, detection a-myc and r-a-mHRP





14/31

Fig.10.





15/31

Fig.11A.



16/31

Fig.11 B.



17/31

Fig.12.





Fig.13.



ApoE 10

Fig.14.





Fig.15.

peptide 1:	biotin-GC-ARMEEMGSRTDRLDE	aa	1-16 (16+2 aa)
peptide 2:	biotin-GC-VKEQVAEVRAKLEEQA	aa	17-32 (16+2 aa)
peptide 3:	biotin-GC-QQIRLQAEAFQARLKS	aa	33-48 (16+2 aa)
peptide 4:	biotin-GC-WFEPLVEDMQRQWAGL	aa	49-64 (16+2 aa)
peptide 5:	biotin-GC-VEKVQAAVGTSAAPVP	aa	65-80 (16+2 aa)
peptide 6:	biotin-GC-RTRDRLDEVKEQVAEV	aa	9-24 (16+2 aa)
peptide 7:	biotin-GC-RAKLEEQAQQIRLQAE	aa	25-40 (16+2 aa)
peptide 8:	biotin-GC-AFQARLKSWFEPLVED	aa	41-56 (16+2 aa)
peptide 9:	biotin-GC-MQRQWAGLVEKVQAAV	aa	57-72 (16+2 aa)
peptide 10:	biotin-GC-GTSAAPVPSDNH	aa	73-84 (12+2 aa)

Fig.16.






Fig.18.




Fig.19.




Fig.20.





Fig.21.



26/31

Fig.22.





Fig.23.



28/31

Fig.24A.




Fig.24B.




Fig.25.




Fig.26.




Fig.27.

